


# MabThera® 1400 mg Comparison Card for SC formulation

Please make sure that you select the correct MabThera® formulation & strength

## SUBCUTANEOUS INJECTION For use in non-Hodgkin's lymphoma ONLY\*†

MabThera 1,400 mg solution for subcutaneous injection




Withdraw directly from the vial and administer by subcutaneous injection 

## INTRAVENOUS INFUSION For use in all MabThera-approved indications†

MabThera 100 mg concentrate for solution for infusion  
MabThera 500 mg concentrate for solution for infusion



Dilute with 0.9% NaCl or 5% glucose and administer by intravenous infusion 

### Check for the specific MabThera SC packaging characteristics before use:

1. Red labelling: 'For subcutaneous use only', 'solution for subcutaneous injection' and 'SC'
2. Pink flip-off cap

\* MabThera SC is not indicated as monotherapy in patients with stage III-IV follicular lymphoma who are chemoresistant or are in their second or subsequent relapse after chemotherapy.

† Please refer to the Summary of Product Characteristics for further information.

### Reporting of side effects

Report any side effects at Roche Bangladesh Local Safety Unit by emailing at [bangladesh.drugsafety@roche.com](mailto:bangladesh.drugsafety@roche.com) or calling +88 01766686086