

# Tarceva®

Erlotinib

---

## 1. DESCRIPTION

### 1.1 Therapeutic/Pharmacologic Class of Drug

Antineoplastic agent.

ATC code L01XE03.

### 1.2 Type of Dosage Form

Tarceva film-coated tablets 25 mg, 100 mg, 150 mg.

Film-coated white to yellowish, round, biconvex tablets, print coloration according to tablet strength.

Or film-coated white to yellowish, round, biconvex tablets, with “T 25 or T 100 or T 150” engraved on one side according to tablet strength.

### 1.3 Route of Administration

Oral.

### 1.4 Sterile/Radioactive Statement

Not applicable.

### 1.5 Qualitative and Quantitative Composition

Active ingredient: erlotinib hydrochloride.

One film-coated tablet of each strength contains erlotinib hydrochloride, corresponding to 25 mg, 100 mg and 150 mg of erlotinib.

#### List of Excipients

##### Tablet core

Lactose monohydrate	Ph. Eur./USP/JP
Cellulose, microcrystalline	Ph. Eur./USP/JP
Sodium starch glycolate	Ph. Eur./USP/JPE
Sodium laurilsulfate	Ph. Eur./USP/JP
Magnesium stearate	Ph. Eur./USP/JP

##### Tablet coat

Hydroxypropyl cellulose	Ph. Eur./USP/JP
Titanium dioxide	Ph. Eur./USP/JP
Polyethylene glycol	Ph. Eur./NF/JP

Hydroxypropyl methylcellulose Ph. Eur./USP/JP

## **2. CLINICAL PARTICULARS**

### **2.1 Therapeutic Indications**

*Non-small cell lung cancer:*

Tarceva is indicated for the first-line and maintenance treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with EGFR-activating mutations.

Tarceva is also indicated for the treatment of patients with locally advanced or metastatic non-small cell lung cancer after failure of at least one prior chemotherapy regimen.

*Pancreatic cancer:*

Tarceva in combination with gemcitabine is indicated for the first-line treatment of patients with locally advanced, unresectable or metastatic pancreatic cancer.

### **2.2 Dosage and Administration**

#### **General**

*Non-small cell lung cancer:*

EGFR mutation testing should be performed prior to initiation of Tarceva as first-line or maintenance therapy in chemo-naïve patients with advanced or metastatic NSCLC.

The recommended daily dose of Tarceva is 150 mg taken at least one hour before or two hours after the ingestion of food.

*Pancreatic cancer:*

The recommended daily dose of Tarceva is 100 mg taken at least one hour before or two hours after the ingestion of food, in combination with gemcitabine (see the label of gemcitabine for the pancreatic cancer indication).

#### **2.2.1 Special Dosage Instructions**

**Drug Interactions:** Concomitant use of CYP3A4 substrates and modulators may require dose adjustment (see section 2.7 Interactions with other Medicinal Products and other Forms of Interaction).

When dose adjustment is necessary, it is recommended to reduce in 50 mg steps (see sections 2.4 Warnings and Precautions and 2.7 Interactions with other Medicinal Products and other Forms of Interaction).

**Hepatic impairment:** Erlotinib is eliminated by hepatic metabolism and biliary excretion. Although erlotinib exposure was similar in patients with moderately impaired hepatic function (Child-Pugh score 7-9) compared with patients with adequate hepatic function, caution should be used when administering Tarceva to patients with hepatic impairment.

Dose reduction or interruption of Tarceva should be considered if severe adverse reactions occur. Safety and efficacy have not been studied in patients with severe hepatic impairment (see sections 2.4.1 General [Warnings and Precautions], *Hepatitis*, *hepatic failure*, and 3.2.5 Pharmacokinetics in Special Populations).

**Renal impairment:** The safety and efficacy of Tarceva has not been studied in patients with renal impairment (see section 3.2.5 Pharmacokinetics in Special Populations).

**Pediatric use:** The safety and efficacy of Tarceva in the approved indications has not been established in patients under the age of 18 years.

**Smokers:** Cigarette smoking has been shown to reduce erlotinib exposure by 50-60%. The maximum tolerated dose of Tarceva in NSCLC patients who currently smoke cigarettes was 300 mg. The 300 mg dose did not show improved efficacy in second-line treatment after failure of chemotherapy compared to the recommended 150 mg dose in patients who continue to smoke cigarettes (see sections 2.7 Interactions with other Medicinal Products and other Forms of Interaction and 3.2.5 Pharmacokinetics in Special Populations).

## 2.3 Contraindications

Tarceva is contraindicated in patients with severe hypersensitivity to erlotinib or to any component of Tarceva.

## 2.4 Warnings and Precautions

### 2.4.1 General

**Interstitial lung disease:** Cases of interstitial lung disease (ILD)-like events, including fatalities, have been reported uncommonly in patients receiving Tarceva for treatment of NSCLC (non-small cell lung carcinoma), pancreatic cancer or other advanced solid tumors.

In pivotal study BR.21, in NSCLC, the incidence of serious ILD-like events was 0.8% in each of the placebo and Tarceva arms.

In a meta-analysis of NSCLC randomized controlled clinical trials, the incidence of ILD-like events was 0.9% on Tarceva compared to 0.4% in patients in the control arms.

In the pancreatic cancer study in combination with gemcitabine, the incidence of ILD-like events was 2.5% in the Tarceva plus gemcitabine group versus 0.4% in the placebo plus gemcitabine-treated group.

Some examples of reported diagnoses in patients suspected of having ILD-like events included pneumonitis, radiation pneumonitis, hypersensitivity pneumonitis, interstitial pneumonia, interstitial lung disease, obliterative bronchiolitis, pulmonary fibrosis, acute respiratory distress syndrome, lung infiltration and alveolitis. These ILD-like events started from a few days to several months after initiating Tarceva therapy. Most of the cases were associated with confounding or contributing factors such as concomitant or prior chemotherapy, prior radiotherapy, pre-existing parenchymal lung disease, metastatic lung disease, or pulmonary infections.

In patients who develop acute onset of new and/or progressive unexplained pulmonary symptoms, such as dyspnea, cough and fever, Tarceva therapy should be interrupted pending diagnostic evaluation. If ILD is diagnosed, Tarceva should be discontinued and appropriate treatment initiated as necessary (see section 2.6 Undesirable Effects).

*Diarrhea, dehydration, electrolyte imbalance and renal failure:* Diarrhea has occurred in patients on Tarceva and moderate or severe diarrhea should be treated with loperamide. In some cases, dose reduction may be necessary. In the event of severe or persistent diarrhea, nausea, anorexia, or vomiting associated with dehydration, Tarceva therapy should be interrupted and appropriate measures should be taken to treat the dehydration (see section 2.6 Undesirable Effects). There have been rare reports of hypokalemia and renal failure (including fatalities). Some reports of renal failure were secondary to severe dehydration due to diarrhea, vomiting and/or anorexia while others were confounded by concomitant chemotherapy. In more severe or persistent cases of diarrhea, or cases leading to dehydration, particularly in groups of patients with aggravating risk factors (concomitant medications, symptoms or diseases or other predisposing conditions including advanced age), Tarceva therapy should be interrupted and appropriate measures should be taken to intensively rehydrate the patients intravenously. In addition, renal function and serum electrolytes including potassium should be monitored in patients at risk of dehydration.

*Hepatitis, hepatic failure:* Rare cases of hepatic failure (including fatalities) have been reported during use of Tarceva. Confounding factors have included pre-existing liver disease or concomitant hepatotoxic medications. Therefore, in such patients, periodic liver function testing should be considered. Tarceva dosing should be interrupted if changes in liver function are severe (see section 2.6 Undesirable Effects).

*Gastrointestinal perforation:* Patients receiving Tarceva are at increased risk of developing gastrointestinal perforation, which was observed uncommonly (including some cases with a fatal outcome). Patients receiving concomitant anti-angiogenic agents, corticosteroids, NSAIDs, and/or taxane-based chemotherapy, or who have prior history of peptic ulceration or diverticular disease are at increased risk. Tarceva should be permanently discontinued in patients who develop gastrointestinal perforation (see section 2.6 Undesirable Effects).

*Bullous and exfoliative skin disorders:* Bullous, blistering and exfoliative skin conditions have been reported, including very rare cases suggestive of Stevens-Johnson syndrome/toxic epidermal necrolysis, which in some cases were fatal (see section 2.6 Undesirable Effects). Tarceva treatment should be interrupted or discontinued if the patient develops severe bullous, blistering or exfoliating conditions.

*Ocular disorders:* Very rare cases of corneal perforation or ulceration have been reported during use of Tarceva. Other ocular disorders including abnormal eyelash growth, keratoconjunctivitis sicca or keratitis have been observed with Tarceva treatment which are also risk factors for corneal perforation/ulceration. Tarceva therapy should be interrupted or discontinued if patients present with acute/worsening ocular disorders such as eye pain (see section 2.6 Undesirable Effects).

*Drug Interactions:* Tarceva has a potential for clinically significant drug-drug interactions (see section 2.7 Interactions with other Medicinal Products and other Forms of Interaction).

#### **2.4.2 Drug Abuse and Dependence**

Not applicable.

#### **2.4.3 Ability to Drive and Use Machines**

Erlotinib has no or negligible influence on the ability to drive and use machines.

### **2.5 Use in Special Populations**

#### **2.5.1 Females And Males of Reproductive Potential**

*Contraception: Females:* Adequate contraceptive methods should be used during therapy, and for at least 2 weeks after completing therapy.

#### **2.5.2 Pregnancy**

There are no adequate or well controlled studies in pregnant women using Tarceva. Studies in animals have shown some reproductive toxicity (see sections 3.3.3, Impairment of Fertility and 3.3.4 Reproductive Toxicity). The potential risk for humans is unknown. Women of childbearing potential must be advised to avoid pregnancy while on Tarceva. Treatment should only be continued in pregnant women if the potential benefit to the mother outweighs the risk to the fetus.

#### **2.5.3 Lactation**

It is not known whether erlotinib is excreted in human milk. No studies have been conducted to assess the impact of Tarceva on milk production or its presence in breast milk. As of the potential for harm to the nursing infant is unknown, mothers should be advised against breastfeeding while receiving Tarceva and for at least 2 weeks after the final dose.

#### **2.5.4 Pediatric Use**

The safety and efficacy of Tarceva in the approved indications has not been established in patients under the age of 18 years (see sections 2.2.1, Special Dosage Instructions and 3.2.5 Pharmacokinetics in Special Population).

#### **2.5.5 Geriatric Use**

See section 3.2.5, Pharmacokinetics in Special Population.

#### **2.5.6 Renal Impairment**

See sections 2.2.1, Special Dosage Instructions and 3.2.5, Pharmacokinetics in Special Population.

## 2.5.7 Hepatic Impairment

Erlotinib exposure was similar in patients with moderately impaired hepatic function (Child-Pugh score 7-9) compared with patients with adequate hepatic function including patients with primary liver cancer or hepatic metastases (see section 2.4.1 General [Warnings and Precautions]). Safety and efficacy have not been studied in patients with severe hepatic impairment (see section 2.2.1, Special Dosage Instructions).

## 2.6 Undesirable Effects

### 2.6.1 Clinical Trials

Safety evaluation of Tarceva is based on the data from more than 1500 patients treated with at least one 150 mg dose of Tarceva monotherapy, and more than 300 patients who received Tarceva 100 mg or 150 mg in combination with gemcitabine.

The incidence of adverse drug reactions (ADRs) reported with Tarceva alone or in combination with chemotherapy are summarized in the tables below and are based on data from clinical trials. The listed ADRs were those reported in at least 10% (in the Tarceva group) of patients and occurred more frequently ( $\geq 3\%$ ) in patients treated with Tarceva than in the comparator arm.

Adverse drug reactions from clinical trials (Table 1) are listed by MedDRA system organ class. The corresponding frequency category for each adverse drug reaction is based on the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1000$ ), very rare ( $< 1/10,000$ ).

### **Non-small cell lung cancer - Tarceva administered as monotherapy**

#### *First-line Treatment of Patients with EGFR Mutations*

In an open-label, randomized phase III study, ML 20650 conducted in 154 patients, the safety of Tarceva for first-line treatment of NSCLC patients with EGFR activating mutations was assessed in 75 patients; no new safety signals were observed in these patients.

The most frequent ADRs seen in patients treated with Tarceva in study ML 20650 were rash and diarrhoea (any Grade 80% and 57%, respectively), most were Grade 1/2 in severity and manageable without intervention. Grade 3 rash and diarrhoea occurred in 9% and 4% of patients, respectively. No Grade 4 rash or diarrhoea was observed. Both rash and diarrhoea resulted in discontinuation of Tarceva in 1% of patients. Dose modifications (interruptions or reductions) for rash and diarrhoea were needed in 11% and 7% of patients, respectively.

#### *Maintenance Treatment*

In two other double-blind, randomized, placebo-controlled Phase III studies (BO18192 and BO25460) conducted in a total of 1532 patients with advanced, recurrent or metastatic NSCLC following first-line standard platinum-based chemotherapy, no new safety signals were identified.

The most frequent ADRs seen in patients treated with Tarceva in studies BO18192 and BO25460 were rash (BO18192: all grades 49.2%, grade 3: 6.0%; BO25460: all grades 39.4%, grade 3: 5.0%) and diarrhea (BO18192: all grades 20.3%, grade 3: 1.8%; BO25460: all grades 24.2%, grade 3: 2.5%) (see table 2). No Grade 4 rash or diarrhoea was observed in either study. Rash and diarrhoea resulted in discontinuation of Tarceva in 1% and < 1% of patients, respectively, in study BO18192, while no patient discontinued for rash or diarrhea in BO25460. Dose modifications (interruptions or reductions) for rash and diarrhoea were needed in 8.3% and 3% of patients, respectively, in study BO18192 and 5.6% and 2.8% of patients, respectively, in study BO25460.

#### *Second and Further Line Treatment*

The ADRs in **Table 1** are based on data from a randomized double-blind study (BR.21) conducted in 17 countries, 731 patients with locally advanced or metastatic NSCLC after failure of at least one prior chemotherapy regimen. Patients were randomized 2:1 to receive Tarceva 150 mg or placebo. Study drug was taken orally once daily until disease progression or unacceptable toxicity.

The most frequent ADRs were rash and diarrhea (any grade 75% and 54%, respectively), most were Grade 1/2 in severity and manageable without intervention. Grade 3/4 rash and diarrhea occurred in 9% and 6%, respectively, in patients treated with Tarceva and each resulted in study discontinuation in 1% of patients. Dose reduction for rash and diarrhea was needed in 6% and 1% of patients, respectively. In study BR.21, the median time to onset of rash was 8 days, and the median time to onset of diarrhea was 12 days.

#### **Pancreatic cancer – Tarceva administered concurrently with gemcitabine**

The ADRs listed in table 1 below are based on erlotinib-arm data from a controlled clinical trial (PA.3), 259 patients with pancreatic cancer received Tarceva 100 mg plus gemcitabine compared to 256 patients in the placebo plus gemcitabine arm.

The most frequent ADRs in study PA.3, in pancreatic cancer patients receiving Tarceva 100 mg plus gemcitabine, were fatigue, rash and diarrhea. In the Tarceva plus gemcitabine arm, Grade 3/4 rash and diarrhea were reported in 5% of patients. The median time to onset of rash and diarrhea was 10 days and 15 days, respectively. Rash and diarrhea each resulted in dose reductions in 2% of patients, and resulted in study discontinuation in up to 1% of patients receiving Tarceva plus gemcitabine.

The Tarceva 150 mg plus gemcitabine cohort (23 patients) was associated with a higher rate of certain class-specific adverse reactions including rash and required more frequent dose reduction or interruption.

**Table 1 ADRs occurring in  $\geq 10\%$  of patients in BR.21 (treated with Tarceva) and PA.3 (treated with Tarceva plus gemcitabine) studies and ADRs occurring more frequently ( $\geq 3\%$ ) than placebo in BR.21 (treated with Tarceva) and PA.3 (treated with Tarceva plus gemcitabine) studies**

NCI-CTC Grade MedDRA Preferred Term	Tarceva (BR.21) N=485			Tarceva (PA.3) N=259			Frequency category of highest incidence
	Any Grade %	3 %	4 %	Any Grade %	3 %	4 %	
<i>Infections and infestations</i> Infection*	24	4	0	31	3	<1	very common
<i>Metabolism and nutrition disorders</i> Anorexia Weight decreased	52 --	8 --	1 --	39 --	2 --	0 --	very common very common
<i>Eye disorders</i> Conjunctivitis Keratoconjunctivitis sicca	12 12	<1 0	0 0	-- --	-- --	-- --	very common very common
<i>Psychiatric disorders</i> Depression	--	--	--	19	2	0	very common
<i>Nervous system disorders</i> Headache Neuropathy	-- --	-- --	-- --	15 13	<1 1	0 <1	very common very common
<i>Respiratory, thoracic and mediastinal disorders</i> Dyspnea Cough	41 33	17 4	11 0	-- 16	-- 0	-- 0	very common very common
<i>Gastrointestinal disorders</i> Diarrhea Nausea Vomiting Stomatitis Abdominal pain Dyspepsia Flatulence	54 33 23 17 11 -- --	6 3 2 <1 2 -- --	<1 0 <1 0 <1 -- --	48 -- -- 22 -- 17 13	5 -- -- <1 -- <1 0	<1 -- -- 0 -- 0 0	very common very common very common very common very common very common very common
<i>Skin and subcutaneous tissue disorders</i> Rash Pruritus Dry skin Alopecia	75 13 12 --	8 <1 0 --	<1 0 0 --	69 -- -- 14	5 -- -- 0	0 -- -- 0	very common very common very common very common

<i>General disorders and administration site conditions</i>	52	14	4	73	14	2	very common
Fatigue	--	--	--	36	3	0	very common
Pyrexia	--	--	--	12	0	0	very common
Rigors							

\* Severe infections, with or without neutropenia, have included pneumonia, sepsis, and cellulitis.

-- corresponds to percentage below threshold

### **Further information on ADRs of special interest**

The following ADRs have been observed in patients who received Tarceva 150 mg as monotherapy or 100 mg or 150 mg in combination with gemcitabine.

Very common ADRs are presented in **Table 1**, ADRs in other frequency categories are summarized below.

#### *Gastrointestinal disorders:*

Gastrointestinal perforations have been reported uncommonly (in less than 1% of patients) with Tarceva treatment, in some cases with a fatal outcome (see section 2.4 Warnings and Precautions).

Cases of gastrointestinal bleeding have been reported commonly (including some fatalities), some associated with concomitant warfarin administration (see also section 2.7 Interactions with other Medicinal Products and other Forms of Interaction), and some with concomitant NSAIDs.

#### *Hepatobiliary disorders:*

Liver function test abnormalities (including raised ALT, AST, bilirubin) have been observed commonly in clinical trials of Tarceva. In study PA.3, these occurred very commonly. They were mainly mild or moderate in severity, transient in nature or associated with liver metastases. Rare cases of hepatic failure (including fatalities) have been reported during use of Tarceva. Confounding factors have included pre-existing liver disease or concomitant hepatotoxic medications (see section 2.4 Warnings and Precautions).

#### *Eye disorders:*

Corneal ulcerations or perforations have been reported very rarely in patients receiving Tarceva treatment (see section 2.4 Warnings and Precautions). Keratitis and conjunctivitis have been reported commonly with Tarceva.

Abnormal eyelash growth, including in-growing eyelashes, excessive growth and thickening of the eyelashes have been reported (see section 2.4 Warnings and Precautions).

#### *Respiratory, thoracic and mediastinal disorders:*

There have been uncommon reports of serious interstitial lung disease (ILD)-like events, (including fatalities), in patients receiving Tarceva for treatment of NSCLC and other advanced solid tumors (see section 2.4 Warnings and Precautions).

Cases of epistaxis have also been reported commonly in both the NSCLC and the pancreatic cancer trials.

*Skin and subcutaneous tissue disorders:*

Rash has been reported very commonly in patients receiving Tarceva and in general, manifests as a mild or moderate erythematous and papulopustular rash, which may occur or worsen in sun-exposed areas. For patients who are exposed to sun, protective clothing, and/or use of sun screen (e.g. mineral-containing) may be advisable. Acne, dermatitis acneiform and folliculitis have been observed commonly, most of these events were mild or moderate and non-serious. Skin fissures, mostly non-serious, were reported commonly and in the majority of cases were associated with rash and dry skin. Other mild skin reactions such as hyperpigmentation have been observed uncommonly (in less than 1% of patients).

Bullous, blistering and exfoliative skin conditions have been reported, including very rare cases suggestive of Stevens-Johnson syndrome/toxic epidermal necrolysis, which in some cases were fatal (see section 2.4 Warnings and Precautions).

Hair and nail changes, mostly non-serious, were reported in clinical trials, e.g. paronychia was reported commonly and hirsutism, eyelash/eyebrow changes and brittle and loose nails were reported uncommonly.

**2.6.2 Postmarketing Experience**

The following adverse drug reactions have been identified from postmarketing experience with Tarceva based on spontaneous case reports and literature cases.

**Table 2 Adverse drug reactions from postmarketing experience**

<b>Adverse reactions</b>	<b>Frequency category</b>
<i>Eye disorders</i>	
Uveitis	Unknown
<i>Skin and subcutaneous tissue disorders</i>	
Hair and nail changes, mostly non-serious, e.g. hirsutism, eyelash/eyebrow changes, paronychia and brittle and loose nails	Uncommon

## 2.7 Interactions with other Medicinal Products and other Forms of Interaction

Erlotinib is metabolized in the liver by the hepatic cytochromes in humans, primarily CYP3A4 and to a lesser extent by CYP1A2, and the pulmonary isoform CYP1A1. Potential interactions may occur with drugs which are metabolized by, or are inhibitors or inducers of, these enzymes.

Potent inhibitors of CYP3A4 activity decrease erlotinib metabolism and increase erlotinib plasma concentrations. Inhibition of CYP3A4 metabolism by ketoconazole (200 mg po BID for 5 days) resulted in increased exposure to erlotinib (86% in median erlotinib exposure [AUC]) and a 69% increase in  $C_{max}$  when compared to erlotinib alone. When Tarceva was co-administered with ciprofloxacin, an inhibitor of both CYP3A4 and CYP1A2, the erlotinib exposure [AUC] and maximum concentration ( $C_{max}$ ) increased by 39% and 17%, respectively. Therefore caution should be used when administering Tarceva with potent CYP3A4 or combined CYP3A4/CYP1A2 inhibitors. In these situations, the dose of Tarceva should be reduced if toxicity is observed.

Potent inducers of CYP3A4 activity increase erlotinib metabolism and significantly decrease erlotinib plasma concentrations. Induction of CYP3A4 metabolism by rifampicin (600 mg p.o. QD for 7 days) resulted in a 69% decrease in the median erlotinib AUC, following a 150 mg dose of Tarceva as compared to Tarceva alone.

Pre-treatment and co-administration of rifampicin with a single 450 mg dose of Tarceva resulted in a mean erlotinib exposure [AUC] of 57.5% of that after a single 150 mg Tarceva dose in the absence of rifampicin treatment. Alternative treatments lacking potent CYP3A4 inducing activity should be considered when possible. For patients who require concomitant treatment with Tarceva and a potent CYP3A4 inducer such as rifampicin an increase in dose to 300 mg should be considered while their safety (see section 2.4.1, Warnings and Precautions, General) is closely monitored, and if well tolerated for more than 2 weeks, further increase to 450 mg could be considered with close safety monitoring. Higher doses have not been studied in this setting.

Pre-treatment or co-administration of Tarceva did not alter the clearance of the prototypical CYP3A4 substrates midazolam and erythromycin. Significant interactions with the clearance of other CYP3A4 substrates are therefore unlikely. Oral availability of midazolam did appear to decrease by up to 24%, which was however not attributed to effects on CYP3A4 activity.

The solubility of erlotinib is pH dependent. Erlotinib solubility decreases as pH increases. Drugs that alter the pH of the upper GI tract may alter the solubility of erlotinib and hence its bioavailability. Co-administration of Tarceva with omeprazole, a proton pump inhibitor, decreased the erlotinib exposure [AUC] and  $C_{max}$  by 46% and 61%, respectively. There was no change to  $T_{max}$  or half-life. Concomitant administration of Tarceva with 300 mg ranitidine, an  $H_2$ -receptor antagonist, decreased erlotinib exposure [AUC] and  $C_{max}$  by 33% and 54%, respectively. Therefore, co-administration of drugs reducing gastric acid production with Tarceva should be avoided where possible. Increasing the dose of Tarceva when co-administered with such agents is not likely to compensate for this loss of exposure. However, when Tarceva was dosed in a staggered

manner 2 hours before or 10 hours after ranitidine 150 mg b.i.d., erlotinib exposure [AUC] and  $C_{max}$  decreased only by 15% and 17%, respectively. If patients need to be treated with such drugs, then an H<sub>2</sub>-receptor antagonist such as ranitidine should be considered and used in a staggered manner. Tarceva must be taken at least 2 hours before or 10 hours after the H<sub>2</sub>-receptor antagonist dosing.

Interaction with coumarin-derived anticoagulants, including warfarin, leading to increased International Normalized Ratio (INR) and bleeding events, which in some cases were fatal, have been reported in patients receiving Tarceva. Patients taking coumarin-derived anticoagulants should be monitored regularly for any changes in prothrombin time or INR.

The combination of Tarceva and a statin may increase the potential for statin-induced myopathy, including rhabdomyolysis, which was observed rarely.

Smokers should be advised to stop smoking as cigarette smoking, which is known to induce CYP1A1 and CYP1A2, has been shown to reduce erlotinib exposure by 50-60%. (See sections, 2.2.1, Special Dosage Instructions, 3.2.5, Pharmacokinetics in Special Populations).

In a phase Ib study, there were no significant effects of gemcitabine on the pharmacokinetics of erlotinib nor were there significant effects of erlotinib on the pharmacokinetics of gemcitabine.

## **2.8 Overdose**

Single oral doses of Tarceva up to 1000 mg in healthy subjects, and up to 1600 mg given as a single dose once weekly in cancer patients, have been tolerated. Repeated twice-daily doses of 200 mg in healthy subjects were poorly tolerated after only a few days of dosing. Based on the data from these studies, severe adverse events such as diarrhea, rash, and possibly liver transaminase elevation may occur above the recommended dose. In case of suspected overdose, Tarceva should be withheld and symptomatic treatment administered.

## **3. PHARMACOLOGICAL PROPERTIES AND EFFECTS**

### **3.1 Pharmacodynamic Properties**

#### **3.1.1 Mechanism of Action**

Erlotinib potently inhibits the intracellular phosphorylation of HER1/EGFR receptor. HER1/EGFR receptor is expressed on the cell surface of normal cells and cancer cells. In non-clinical models, inhibition of EGFR phosphotyrosine results in cell stasis and/or death.

#### **3.1.2 Clinical/Efficacy Studies**

*Non-small cell lung cancer (Tarceva administered as monotherapy):*

*First-line therapy for patients with EGFR-activating mutations:*

The efficacy of Tarceva in first-line treatment of patients with EGFR-activating mutations in NSCLC was demonstrated in a phase III, randomized, open-label trial (ML20650, EURTAC). This study was conducted in Caucasian patients with metastatic or locally advanced NSCLC (stage IIIB and IV) who have not received previous chemotherapy or any systemic antitumor therapy for their advanced disease and who present mutations in the tyrosine kinase domain of the EGFR (exon 19 deletion or exon 21 mutation). Patients were randomized 1:1 to receive Tarceva 150 mg or platinum-based doublet chemotherapy.

The primary endpoint of investigator-assessed progression-free survival (PFS) was determined at a preplanned interim analysis (n=153, hazard ratio [HR] = 0.42, 95 % CI, 0.27 to 0.64; p<0.0001 for the Tarceva group [n=77] relative to the chemotherapy group [n=76]). A 58% reduction in the risk of disease progression or death was observed. In the Tarceva versus chemotherapy arms, respectively, median PFS was 9.4 and 5.2 months and objective response rate (ORR) was 54.5% and 10.5%. PFS results were confirmed by an independent review of the scans, median PFS was 10.4 months in the Tarceva group compared with 5.4 months in the chemotherapy group (HR=0.47, 95% CI, 0.27 to 0.78; p=0.003). The overall survival data were immature at the time of interim analysis (HR=0.80, 95% CI, 0.47 to 1.37; p=0.4170).

At an updated analysis with 62% of OS maturity, OS HR was 0.93 (95% CI, 0.64 to 1.36; p=0.7149). A high crossover was observed with 82% of the patients in the chemotherapy arm receiving subsequent therapy with an EGFR tyrosine kinase inhibitor and all but 2 of those patients had subsequent Tarceva. In the updated analysis, PFS results remained consistent with the interim analysis results. Median PFS assessed by the investigators was 10.4 and 5.1 months in the Tarceva and chemotherapy arms, respectively (HR=0.34, 95% CI, 0.23 to 0.49; p<0.0001).

- *Additional published data*

In a prospective analysis of patients with advanced NSCLC having tumors with activating mutations in the EGFR TK domain, the median PFS for the 113 patients treated with first-line Tarceva was 14 months (95% CI, 9.7 to 18.3 months) and the median overall survival was 28.0 months (95% CI, 22.7 to 33 months).

A pooled analysis of published data from NSCLC patients showed that patients having tumors with EGFR-activating mutations and receiving Tarceva as predominantly first-line therapy (n=70, 12.5 months, 95% CI [10.6-16.0]) had a longer median PFS compared to those receiving chemotherapy (n=359, 6.0 months, 95% CI [5.4-6.7]).

*First-line maintenance therapy:*

The efficacy and safety of Tarceva as first-line maintenance therapy of NSCLC was investigated in a randomized, double-blind, placebo-controlled trial BO18192 (SATURN). This study was conducted in 889 patients with locally advanced or metastatic NSCLC who did not progress during 4 cycles of platinum-based doublet chemotherapy. Patients were randomized 1:1 to receive Tarceva 150 mg or placebo orally once daily. The primary endpoint of the study was PFS in all patients and in patients with

an EGFR IHC-positive tumor. Baseline demographic and disease characteristics were well balanced between the two treatment arms.

In this study BO18192 (SATURN), the overall population showed a benefit for the primary PFS endpoint (HR=0.71;  $p<0.0001$ ) and the secondary OS endpoint (HR=0.8;  $p=0.0088$ ). However, the largest benefit was observed in a predefined exploratory analysis in patients with EGFR-activating mutations ( $n=49$ ) demonstrating a substantial PFS benefit (HR=0.10, 95% CI, 0.04 to 0.25;  $p<0.0001$ ) and an overall survival HR of 0.83 (95% CI, 0.34 to 2.02). 67% of placebo patients in the EGFR mutation-positive subgroup received second- or further-line treatment with EGFR TKIs. In patients with EGFR wild-type tumors ( $n=388$ ), the PFS HR was 0.78 (95% CI, 0.63 to 0.96;  $p=0.0185$ ) and the overall survival HR was 0.77 (95% CI, 0.61 to 0.97;  $p=0.0243$ ).

The BO25460 (IUNO) study was conducted in 643 patients with advanced NSCLC whose tumors did not harbor an EGFR-activating mutation (exon 19 deletion or exon 21 L858R mutation) and who had not experienced disease progression after four cycles of platinum-based chemotherapy.

The objective of the study was to compare the overall survival of first-line maintenance therapy with erlotinib versus erlotinib administered at the time of disease progression. The study did not meet its primary endpoint. OS of Tarceva in first-line maintenance was not superior to Tarceva as second-line treatment in patients whose tumor did not harbor an EGFR-activating mutation (HR=1.02, 95% CI, 0.85 to 1.22;  $p=0.82$ ). The secondary endpoint of PFS showed no difference between Tarceva and placebo in maintenance treatment (HR=0.94, 95% CI, 0.80 to 1.11;  $p=0.48$ ).

Based on the data from the BO25460 (IUNO) study, Tarceva use is not recommended for first-line maintenance treatment in patients without an EGFR-activating mutation.

#### *Second-/third-line therapy:*

The efficacy and safety of Tarceva in second-/third-line therapy of NSCLC was demonstrated in a randomized, double-blind, placebo-controlled trial (BR.21). This study was conducted in 17 countries, in 731 patients with locally advanced or metastatic NSCLC after failure of at least one chemotherapy regimen. Patients were randomized 2:1 to receive Tarceva 150 mg or placebo orally once daily. Study endpoints included overall survival, time to deterioration of lung cancer-related symptoms (cough, dyspnea and pain), response rate, duration of response, PFS, and safety. The primary endpoint was survival.

Due to the 2:1 randomization, 488 patients were randomized to Tarceva and 243 patients to placebo. Patients were not selected for HER1/EGFR status, gender, race, smoking history and histological classification.

Demographic characteristics were well balanced between the two treatment arms. About two-thirds of the patients were male and approximately one-third had a baseline ECOG performance status (PS) of 2 and 9% had a baseline ECOG of 3. Ninety-three percent and 92% of all patients in the Tarceva and placebo groups, respectively, had received a prior platinum-containing regimen and 36% and 37% of all patients, respectively, had received

a prior taxane therapy. Fifty percent of the patients had received only one prior regimen of chemotherapy.

Survival was evaluated in the intent-to-treat population. The median overall survival improved by 42.5% and was 6.7 months in the Tarceva group (95% CI, 5.5 to 7.8 months) compared with 4.7 months in the placebo group (95% CI, 4.1 to 6.3 months). The primary survival analysis was adjusted for the stratification factors as reported at the time of randomization (ECOG PS, best response to prior therapy, number of prior regimens, and exposure to prior platinum) and HER1/EGFR status. In this primary analysis, the adjusted hazard ratio for death in the Tarceva group relative to the placebo group was 0.73 (95% CI, 0.60 to 0.87) ( $p=0.001$ ). The percentage of patients alive at 12 months was 31.2% and 21.5%, respectively.

The survival benefit with Tarceva treatment was seen across most subsets. A series of subsets of patients formed by the values of the stratification factors at randomization and at baseline, HER1/EGFR status, prior exposure to taxanes, smoking history, gender, age, histology, prior weight loss, time between initial diagnosis and randomization, and geographic location were examined in exploratory univariate analyses to assess the robustness of the overall survival result. Nearly all of the hazard ratios in the Tarceva group relative to the placebo group were less than 1.0, suggesting that the survival benefit from Tarceva was robust across subsets. Of note, the survival benefit of Tarceva was comparable in patients with a baseline ECOG PS of 2-3 (HR=0.77) or a PS of 0-1 (HR=0.73) and patients who had received one chemotherapy regimen (HR=0.76) or two or more regimens (HR=0.76).

A survival benefit of Tarceva was also observed in patients who did not achieve an objective tumor response (by RECIST). This was evidenced by a hazard ratio for death of 0.83 among patients whose best response was stable disease and 0.85 among patients whose best response was progressive disease.

**Table 3** summarizes the results for the study, including survival, time to deterioration of lung cancer-related symptoms, and progression-free survival (PFS).

**Table 3: Study BR.21 efficacy results**

	<b>Tarceva (n=488)</b>	<b>Placebo (n=243)</b>	<b>P-value</b>
Median survival 95% CI	6.7 months (5.5 to 7.8)	4.7 months (4.1 to 6.3)	
Difference between survival curves			0.001
Hazard ratio*, mortality (erlotinib:placebo) 95% CI (risk ratio)	0.73 0.60 to 0.87		
Median time to deterioration in cough*** 95% CI	28.1 weeks (16.1 to 40.0)	15.7 weeks (9.3 to 24.3)	0.041
Median time to deterioration in dyspnea*** 95% CI	20.4 weeks (16.3 to 28.3)	12.1 weeks (9.3 to 20.9)	0.031**
Median time to deterioration in pain*** 95% CI	12.1 weeks (10.1 to 14.1)	8.1 weeks (7.7 to 12.3)	0.040**
Median progression-free survival 95% CI	9.7 weeks (8.4 to 12.4)	8.0 weeks (7.9 to 8.0)	<0.001

\* Adjusted for stratification factors and HER1/EGFR status; a value less than 1.00 favors Tarceva (primary analysis).

\*\* P-value adjusted for multiple testing.

\*\*\* From the EORTC QLQ-C30 and QLQ-LC13 Quality of Life Questionnaire.

Symptom deterioration was measured using the EORTC QLQ-C30 and QLQ-LC13 quality of life questionnaires. Baseline scores of cough, dyspnea and pain were similar in the two treatment groups. Tarceva resulted in symptom benefits by significantly prolonging time to deterioration in cough (HR=0.75), dyspnea (HR=0.72) and pain (HR=0.77) versus placebo. These symptom benefits were not due to an increased use of palliative radiotherapy or concomitant medication in the Tarceva group.

The median PFS was 9.7 weeks in the Tarceva group (95% CI, 8.4 to 12.4 weeks) compared with 8.0 weeks in the placebo group (95% CI, 7.9 to 8.1 weeks). The HR for progression, adjusted for stratification factors and HER1/EGFR status, was 0.61 (95% CI, 0.51 to 0.73) (p<0.001). The percent of PFS at 6 months was 24.5% and 9.3%, respectively, for the Tarceva and placebo arms.

The objective response rate by RECIST in the Tarceva group was 8.9% (95% CI, 6.4 to 12.0%). The median duration of response was 34.3 weeks, ranging from 9.7 to 57.6+ weeks. Two responses (0.9%, 95% CI, 0.1 to 3.4) were reported in the placebo group. The proportion of patients who experienced complete response, partial response or stable disease was 44.0% and 27.5%, respectively, for the Tarceva and placebo groups (p=0.004).

In a double-blind, randomized phase III study (MO22162, CURRENTS) comparing two doses of Tarceva (300 mg versus 150 mg) in current smokers (mean of 38 pack years) with locally advanced or metastatic NSCLC in the second-line setting after failure on chemotherapy, the 300 mg dose of Tarceva demonstrated no PFS benefit over the recommended dose (7.00 vs 6.86 weeks, respectively). Patients in this study were not selected based on EGFR mutation status.

*Pancreatic cancer (Tarceva administered concurrently with gemcitabine):*

The efficacy and safety of Tarceva in combination with gemcitabine as a first-line treatment was assessed in a randomized, double-blind, placebo-controlled trial in 569 patients with locally advanced, unresectable or metastatic pancreatic cancer. Patients were randomized 1:1 to receive Tarceva (100 mg or 150 mg) or placebo once daily on a continuous schedule plus gemcitabine i.v. (1000 mg/m<sup>2</sup>, cycle 1 - days 1, 8, 15, 22, 29, 36 and 43 of an 8-week cycle; cycle 2 and subsequent cycles - days 1, 8 and 15 of a 4-week cycle [approved dose and schedule for pancreatic cancer, see the gemcitabine SPC]). Tarceva or placebo was taken orally once daily until disease progression or unacceptable toxicity. Study endpoints included overall survival, response rate and PFS. Duration of response was also examined. The primary endpoint was survival. A total of 285 patients were randomized to receive gemcitabine plus Tarceva (261 patients in the 100 mg cohort and 24 patients in the 150 mg cohort) and 284 patients were randomized to receive gemcitabine plus placebo (260 patients in the 100 mg cohort and 24 patients in the 150 mg cohort). Too few observations were made for the 150 mg cohort to draw conclusions.

Baseline demographic and disease characteristics of the patients were similar between the 2 treatment groups, 100 mg Tarceva plus gemcitabine or placebo plus gemcitabine, except for a slightly larger proportion of females in the Tarceva arm (51%) compared with the placebo arm (44%). The median time from initial diagnosis to randomization was approximately 1.0 month. Approximately half of the patients had a baseline ECOG performance status (PS) of 1, and 17% had a baseline ECOG PS of 2. Most patients presented with metastatic disease at study entry as the initial manifestation of pancreatic cancer (77% in the Tarceva arm, 76% in the placebo arm).

Survival was evaluated in the intent-to-treat population based on follow-up survival data including 551 deaths. Results are presented for the 100 mg dose cohort (504 deaths). The adjusted hazard ratio for death in the Tarceva group relative to the placebo group was 0.82 (95% CI, 0.69 to 0.98) (p=0.028). The percent of patients alive at 12 months was 23.8% in the Tarceva group compared to 19.4% in the placebo group. The median overall survival was 6.4 months in the Tarceva group compared with 6 months in the placebo group.

**Table 4** summarizes the results of the study.

**Table 4: Study PA.3 efficacy results**

	<b>Tarceva 100 mg plus gemcitabine (n=261)</b>	<b>Placebo plus gemcitabine (n=260)</b>	<b>P-value</b>
Median survival	6.4 months	6 months	
Hazard ratio, mortality (erlotinib:placebo) (95% CI)	0.82 (0.69 to 0.98)		p=0.028
% patients alive at 12 months	23.8	19.4	

The median PFS was 3.81 months (16.5 weeks) in the Tarceva group (95% CI, 3.58 to 4.93 months) compared with 3.55 months (15.2 weeks) in the placebo group (95% CI, 3.29 to 3.75 months) (p=0.006).

The median duration of response was 23.9 weeks, ranging from 3.71 to 56+ weeks. The objective response rate (complete response and partial response) was 8.6% in the Tarceva group and 7.9% in the placebo group. The proportion of patients who experienced complete response, partial response or stable disease was 59% and 49.4%, respectively, for the Tarceva and placebo groups (p=0.036).

### **3.1.3 Immunogenicity**

Not applicable.

## **3.2 Pharmacokinetic Properties**

### *Exposure:*

Following a 150 mg oral dose of Tarceva, at steady state, the median time to reach maximum plasma concentrations is approximately 4.0 hours with median maximum plasma concentrations achieved of 1,995 ng/ml. Prior to the next dose at 24 hours, the median minimum plasma concentrations are 1,238 ng/ml. Median AUC achieved during the dosing interval at steady state is 41.300 mcg·hr/ml.

### **3.2.1 Absorption**

Oral erlotinib is well absorbed and has an extended absorption phase, with mean peak plasma levels occurring at 4 hours after oral dosing. A study in normal healthy volunteers provided an estimate of bioavailability of 59%. The exposure after an oral dose may be increased by food.

Following absorption, erlotinib is highly bound in blood, with approximately 95% bound to blood components, primarily to plasma proteins (i.e. albumin and alpha-1 acid glycoprotein [AAG]), with a free fraction of approximately 5%.

### **3.2.2 Distribution**

Erlotinib has a mean apparent volume of distribution of 232 l and distributes into tumor tissue of humans. In a study of 4 patients (3 with non-small cell lung cancer [NSCLC], and 1 with laryngeal cancer) receiving 150 mg daily oral doses of Tarceva, tumor samples from surgical excisions on day 9 of treatment revealed tumor concentrations of erlotinib that averaged 1,185 ng/g of tissue. This corresponded to an overall average of 63% of the observed steady-state peak plasma concentrations. The primary active metabolites were present in the tumor at concentrations averaging 160 ng/g tissue, which corresponded to an overall average of 113% of the observed steady-state peak plasma concentrations. Tissue distribution studies using whole body autoradiography following oral administration with [<sup>14</sup>C]-labeled erlotinib in athymic nude mice with HN5 tumor xenografts have shown rapid and extensive tissue distribution with maximum

concentrations of radiolabeled drug (approximately 73% of that in plasma) observed at 1 hour.

### 3.2.3 Metabolism

Erlotinib is metabolized in humans by hepatic cytochrome P450 enzymes, primarily by CYP3A4 and to a lesser extent by CYP1A2. Extrahepatic metabolism by CYP3A4 in intestine, CYP1A1 in lung, and CYP1B1 in tumor tissue potentially contribute to the metabolic clearance of erlotinib. *In vitro* studies indicate approximately 80-95% of erlotinib metabolism by the CYP3A4 enzyme. There are three main metabolic pathways identified: 1) O-demethylation of either side chain or both, followed by oxidation to the carboxylic acids; 2) oxidation of the acetylene moiety followed by hydrolysis to the aryl carboxylic acid; and 3) aromatic hydroxylation of the phenyl-acetylene moiety. The primary metabolites of erlotinib produced by O-demethylation of either side chain have comparable potency to erlotinib in nonclinical *in vitro* assays and *in vivo* tumor models. They are present in plasma at levels that are <10% of erlotinib and display similar pharmacokinetics as erlotinib.

### 3.2.4 Elimination

The metabolites and trace amounts of erlotinib are excreted predominantly via the feces (>90%), with renal elimination accounting for only a small amount of an oral dose.

#### *Clearance:*

A population pharmacokinetic analysis in 591 patients receiving single-agent Tarceva shows a mean apparent clearance of 4.47 l/hr with a median half-life of 36.2 hours. Therefore, the time to reach steady-state plasma concentration would be expected to occur in approximately 7-8 days. No significant relationships between predicted apparent clearance and patient age, body weight, gender, and ethnicity were observed.

Patient factors, which correlate with erlotinib pharmacokinetics, are serum total bilirubin, AAG concentrations and current smoking. Increased serum concentrations of total bilirubin and AAG concentrations were associated with a slower rate of erlotinib clearance. Smokers had a higher rate of erlotinib clearance (see section 2.7 Interactions with other Medicinal Products and other Forms of Interaction).

A second population pharmacokinetic analysis was conducted that incorporated erlotinib data from 204 pancreatic cancer patients who received erlotinib plus gemcitabine. This analysis demonstrated that covariates affecting erlotinib clearance in patients from the pancreatic study were very similar to those seen in the prior single-agent pharmacokinetic analysis. No new covariate effects were identified. Co-administration of gemcitabine had no effect on erlotinib plasma clearance.

### 3.2.5 Pharmacokinetics in Special Populations

There have been no specific studies in pediatric or elderly patients.

Hepatic impairment: Erlotinib is mainly cleared by the liver. Erlotinib exposure was similar in patients with moderately impaired hepatic function (Child-Pugh score 7-9)

compared with patients with adequate hepatic function including patients with primary liver cancer or hepatic metastases.

**Renal impairment:** Erlotinib and its metabolites are not significantly excreted by the kidneys, as less than 9% of a single dose is excreted in the urine. No clinical studies have been conducted in patients with compromised renal function.

**Smokers:** A pharmacokinetic study in non-smoking and currently cigarette-smoking healthy subjects has shown that cigarette smoking leads to increased clearance of, and decreased exposure to, erlotinib. The AUC<sub>0-infinity</sub> in smokers was about 1/3 of that in never/former smokers (n=16 in each of smoker and never/former smoker arms). This reduced exposure in current smokers is presumably due to induction of CYP1A1 in the lung and CYP1A2 in the liver.

In the pivotal phase III NSCLC trial, current smokers achieved erlotinib steady-state trough plasma concentration of 0.65 µg/ml (n=16) which was approximately 2-fold less than the former smokers or patients who had never smoked (1.28 µg/ml, n=108). This effect was accompanied by a 24% increase in apparent erlotinib plasma clearance.

In a phase I dose escalation study in NSCLC patients who were current smokers, pharmacokinetic analyses at steady state indicated a dose-proportional increase in erlotinib exposure when the Tarceva dose was increased from 150 mg to the maximum tolerated dose of 300 mg. Steady-state trough plasma concentrations at a 300 mg dose in current smokers in this study was 1.22 µg/ml (n=17) (see sections 2.2.1 Special Dosage Instructions and 2.7 Interactions with other Medicinal Products and other Forms of Interaction).

### **3.3 Nonclinical Safety**

#### **3.3.1 Carcinogenicity**

Evidence for a carcinogenic potential was not seen in nonclinical studies. Erlotinib was neither genotoxic nor clastogenic in genetic toxicity studies. Two-year carcinogenicity studies with erlotinib conducted in rats and mice at exposures exceeding human therapeutic exposure were negative.

#### **3.3.2 Genotoxicity**

Erlotinib was negative in the standard battery of genotoxicity assays.

#### **3.3.3 Impairment of Fertility**

Impairment of fertility was not observed in studies with male and female rats at doses near the MTD levels.

#### **3.3.4 Reproductive Toxicity**

Data from reproductive toxicology tests in rats and rabbits indicate that, following exposure to erlotinib at doses near the MTD and/or doses that were maternally toxic, there was embryotoxicity, but there was no evidence of teratogenicity, or abnormal pre- or postnatal physical or behavioral development. Maternal toxicity in both rats and

rabbits in these studies occurred at plasma exposure levels that were similar to those in humans following a 150 mg dose of erlotinib.

### **3.3.5 Other**

Chronic dosing effects observed in at least 1 animal species or study included effects on the cornea (atrophy, ulceration), skin (follicular degeneration and inflammation, redness, and alopecia), ovary (atrophy), liver (liver necrosis), kidney (renal papillary necrosis and tubular dilatation), and gastrointestinal tract (delayed gastric emptying and diarrhea). Red blood cell (RBC) counts, hematocrit and hemoglobin were decreased and reticulocytes were increased. White blood cells (WBCs), primarily neutrophils, were increased. There were treatment-related increases in alanine aminotransferase (ALT), aspartate aminotransferase (AST), and bilirubin.

*In vitro* studies of erlotinib showed inhibition of hERG channels at concentrations at least 20 times higher than the free drug concentration in humans at therapeutic doses. Studies in dogs did not show QT prolongation. A systematic centralized review of ECG data from 152 individuals from seven studies with healthy volunteers found no evidence of QT prolongation, and clinical studies have found no evidence of arrhythmias, associated with QT prolongation.

## **4. PHARMACEUTICAL PARTICULARS**

### **4.1 Storage**

#### **Stability**

Tarceva should not be used after the expiry date (EXP) shown on the pack.

Tablets remain stable in their packs for 4 years.

#### **Special Precautions for Storage**

Do not store above 30°C.

### **4.2 Special Instructions for Use, Handling and Disposal**

#### *Disposal of unused/expired medicines*

The release of pharmaceuticals in the environment should be minimized. Medicines should not be disposed of via wastewater and disposal through household waste should be avoided.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

### **4.3 Packs**

25 mg tablets	30
100 mg tablets	30
150 mg tablets	30

Medicine: keep out of reach of children

Current at December 2017

Made for F. Hoffmann-La Roche Ltd, Basel, Switzerland

by Roche S.p.A. Milan, production site Segrate, Italy